Product data sheet

Natupol

Bombus terrestris | Buff-tailed bumblebee



- Standard bumblebee hive for use in protected crops
- With Beevision print
- With Beehome door
- With insulation of brood

Objective

Natupol can be used for a wide range of crops, grown on surfaces bigger than 2000 m² and with approximately 25-35 flowers per m² per week.

Use for

Crops

Round, beef and plum tomatoes and other vegetable or fruit crops.

How it works

Mode of action

After introduction of the hive, worker bumblebees start pollinating the flowers while at the same time collecting pollen to feed the brood. More workers emerge from the brood in the weeks after the introduction, increasing both colony size and pollination performance. After some weeks the colony will reach its maximum size and starts declining in size and pollination activity. Colony development depends on environmental conditions and the amount and quality of pollen.



Bombus terrestris

Product specifications

Presentation
One standard colony of bumblebees including a queen, workers, brood and sugar water.

Colony size
Standard size, containing 60-80 workers at introduction.

Optimum pollination activity
5-7 weeks.

Volume sugar water
1.6 litres.

Insulation
Insulation under brood enabling bumblebees to maintain brood temperature during cold conditions.


Ventilation
Standard ventilation opening.

Beevision print
Equipped with Beevision print, to provide guidance for bumblebees in crops.

Hive door
Equipped with Beehome door to collect bumblebees in cases where bumblebees or hives have to be temporarily removed.

Directions for use

Application
- Place the hive on a horizontal platform (avoid inclination, otherwise sugar water may leak)
- Position the hive about 20-60 cm above the ground in the shade of the crop or if necessary provide additional shading
- Under short day conditions (<10 hours light) position the hive near the top of plants, in such a way that the hive is exposed to the first daylight
- Wait for at least 15 minutes after positioning the hives before opening the hive entrance, this allows the bumblebees to calm down
- Pull the white sliding door of the hive upward until two openings become visible (entrance and exit). Do not remove the small grey door under the lid
- Pollination performance and duration varies per crop and depends on environmental conditions. Always consult a Koppert advisor or a recognized distributor of Koppert products for advice on the best strategy for your situation



Watch the Application video of Natupol

Environmental conditions
Natupol performs best at ambient temperatures between 12-30°C. Pay attention to advice for best practice under conditions with artificial light or in hot environments.

Side Effects
Pesticides can have (in)direct effects on biological solutions. Find out which pesticides have side effects on this product.

Product handling

Storage time after receipt
To guarantee the best pollination results, start using the hive on the day of arrival. If necessary hives can be stored for maximum 1 day.

Storage temperature
15–25°C.

Storage conditions
Keep hives in a well ventilated environment, out of reach of direct sunlight.

Hygiene
- Never move a hive from one greenhouse to another
- Remove bumblebee hives the latest 10 weeks after introduction
- Clean the hive stand before placing new hives

Disposal
Seal old hives into plastic bags for approximately 2 weeks. After this period dispose the hives according to local legislation.

Notes

A number of videos about pollination with Koppert products are available on Koppert's Youtube channel.

Disclaimers

The general conditions of Koppert (Koppert B.V. and/or of its affiliated companies) apply. Only use products that are permitted in your country/state and crop. Always comply with the conditions specified in local product registrations. Koppert cannot be held liable for unauthorized use. Koppert is not liable for any loss of quality if the product is stored for longer than recommended and/or under incorrect conditions.