Prestop WP Compatibility Lists

Fungicides

Active Substance	Recommended Interval (days)	Product Examples
Azoxystrobin	2	Amistar, Ortiva
Benthiavalicarb-isopropyl + Folpet	1	Vincare
Boscalid + Kresoxim-methyl	0	Collis
Boscalid + Pyraclostrobin	2	Signum, Insignis, Bellis, Daisy
Bupirimate	0	Nimrod
Carboxin	4	Vitavax, Krisvax
Chlorothalonil	0	Bravo, Daconil
Copper Oxychloride	0	Cuprokylt
Cyflufenamid	0	Takumi SC, Cyflamid
Cymoxanil	0	Curzate, Option
Dithianon	1	Dithianon WG, Rathianon 70 WG
Dodemorph-acetate	0	Meltatox
Fenhexamid	0	Teldor
Fludioxonil	4	Beret Gold
Fludioxonil + Cyprodinil	4	Switch, Reversal
Fluopyram	0(1)*	Luna Sensation contains this active substance + Trifloxystrobin
Flutriafol	0	Consol, Impact, Pointer
Fosetyl-Aluminium	0	Plant Trust
Hymexazol	7	Tachigaren 70 WP
Imazalil	4	Fungazil 50 LS, Gavel
Iprodione	4	Rovral WG, Rovral Aquaflo, Chipco Green
Kresoxim-methyl	0	Beem, Fermor 500WG, Stroby WG, Strong WG
Magnesium Sulphate + Orange Oil	0	Prev-Magnum
Mancozeb	4	Dithane DF Neotec, Dithane 945, Karamate DF Neotec
Mepanipyrim	0	Frupica SC
Metalaxyl-M	0	Apron XL, Floreo, Fongarid Gold, SL567A, Subdue
Myclobutanil	0	Systhane 20 EW, Crassus
Myclobutanil + Cyclohexanone	0	Clayton Lithium
Myclobutanil + Quinoxyfen	1	Arius
Penconazole	1	Topenco, Topas
Potassium iodide + Potassium thiocyanate + Addit	2**	Enzicur + Addit adjuvant
Prochloraz	7	Sporgon 50 WP, Sportak 45 EW

Prestop WP Compatibility Lists

Active Substance	Recommended Interval (days)	Product Examples
Prochloraz + Propiconazole	7	Bumper P, Cirkon
Procymidone	0	Sumisclex
Propamocarb-hydrochloride	0	Edipro, Promess, Proplant
Propamocarb-hydrocloride + Fosetyl-Aluminium	0	Avatar, Pan Cradle, Previcur Energy
Pyrimethanil	1	Scala, Pyrimala, Pyrus 400SC, Spectrum
Quinoxyfen	0	Apres, Fortress
Sulphur	1	Kumulus DF
Thiophanate-methyl	2	Cercobin, Taurus, Topsin WG
Thiram	4	Thiraflo, Thyram Plus
Tolclofos-methyl	2	Basilex, Rizolex
Trifloxystrobin	0	Flint, Martinet, Swift SC
Trifloxystrobin + Propiconazole	4	Stratego
Triflumizole	0	Procure, Rocket EC, Trifmine

0 (1)* An interval of 1 day is required when PRESTOP® and fluopyram are applied on the same part of the plant. When root system is treated with fluopyram, Prestop can be sprayed in the foliage during the same day.

2 Prestop can be sprayed on the foliage two days after application of Enzicur. Re-application of PRESTOP® is necessary after repeated Enzicur treatments.**

- **The recommended interval between applications of PRESTOP® and a chemical product not mentioned in this list is 7 days.**
- **PRESTOP® can be applied in mixtures with compatible products. However, the mixture should be applied immediately after preparation.**
- **Do not tank mix PRESTOP® with concentrated solutions of pesticides or fertilisers.**

Insecticides

Active Substance	Recommended Interval (days)	Product Examples
Buprofezin	0	Applaud 25 EC, Applaud 25 WP
Cypermethrin	2	Permasect C, Toppel 100
Deltamethrin	0	Bandu, Decis
Imidacloprid	0	Couraze, Imidasect 5GR, Merit Forest
Pyrethrins	0	Pyrethrum 5 EC, Spruzit
Pirimicarb	0	Aphox, Clayton Pirimicarb, Phantom
Spinosad	0	Conserve, Exocet, Tracer
Tebufenozide	0	Mimic, Confirm, Romdan

- **The recommended interval between applications of PRESTOP® and a chemical product not mentioned in this list is 7 days.**
- **PRESTOP® can be applied in mixtures with compatible products. However, the mixture should be applied immediately after preparation.**
- **Do not tank mix PRESTOP® with concentrated solutions of pesticides or fertilisers.**

Prestop WP Compatibility Lists

Biological Control Agents

Active Substance	Recommended Interval (days)	Product Examples
Beneficial arthropods & nematodes	0	Numerous – all are compatible
Bacillus thuringiensis	0	Agree 50 WG, Bruco, Dipel DF, Lepinox Plus, Gnatrol SC, Biocure
Beauveria bassiana	0	Naturalis-L, Botanigard WP
Metarhizium anisopliae	0	Met 52 GR, Met 52 OD
Rape Seed Oil	0	Carbon Kick Booster

Adjuvants

Product Name	Recommended Interval (days)
Addit	0***(<0.25%)
Silwet Gold	0

0* Prestop is compatible with Addit up to the maximum concentration of 0.25%. Higher concentrations of Addit are harmful to Prestop.**

- **The recommended interval between applications of PRESTOP® and a chemical product not mentioned in this list is 7 days.**
- **PRESTOP® can be applied in mixtures with compatible products. However, the mixture should be applied immediately after preparation.**
- **Do not tank mix PRESTOP® with concentrated solutions of pesticides or fertilisers.**

Growth Regulators

Product Name	Recommended Interval (days)
Pirouette	0

Disinfectants

Product Name	Recommended Interval (days)
Resiclean	Compatible (<100 ppm)
Other	1 day

This information was prepared by: Andrew Gough, LPC Technical Manager

Date: 5th July 2017

Key: Product names in red are not on the UK (HSE CRD) databases at the date of compilation